

ARB Compared with non-ARB in Preventing Cardiac Events in High Risk Hypertensive Patients: An Evidence Based Case Report

Alvin Nursalim,* Yoga Yuniadi#

Background: High risk hypertensive patients have an increased risk of developing cardiovascular complication. It is better to use a proven cardioprotective drugs to reduce blood pressure in high risk hypertensive patients. Angiotensin II Receptor Blocker (ARB) is one type of antihypertensive drugs with cardioprotective effect for hypertensive patients without other risk factor. Whether cardioprotective effect of ARB also apply for a more specific population such as high risk hypertensive patients need to be investigated.

Aim: To determine the efficacy of ARB compared to non-ARBs in preventing cardiac event in a more specific population, such as high risk hypertensive patients.

Methods: A search was conducted on PubMed and Cochrane. The selection of title and abstract was done using inclusion and exclusion criterias. Three original articles were found and used as the evidence for the clinical question. The selected studies were critically appraised for validity, importance and applicability.

Result: According to Sawada et al, the blood pressure lowering effect was similar between valsartan and non-ARB groups. The cardiovascular events in valsartan group is lower compared to non-ARB groups (relative risk: 0.54, 95% confidence interval 0.4-0.7, $p < 0.001$). The administration of valsartan as compared to non-ARB, also reduce the occurrence of angina pectoris (Relative risk : 0.52, 95% Confidence Interval 0.31–0.86, $P = 0.01058$). Cohn JN et al showed that there was no significant differences in the candesartan group in terms of total death and primary endpoints. The only significant finding in this article was the lower rate of diabetes mellitus in the candesartan group.

Conclusion: Valsartan, as compared to non-ARB, reduce cardiac event in high risk hypertensive patients.

(J Kardiol Indones. 2012;33:91-8)

Keywords: Angiotensin II receptor blocker, valsartan, high risk hypertension, cardiovascular complication.

*Student of Faculty of Medicine University of Indonesia,

#Departement of Cardiology and Vascular Medicine, Faculty of Medicine, University of Indonesia, and National Cardiovascular Center Harapan Kita, Jakarta

ARB dibandingkan non-ARB dalam Menurunkan Komplikasi Kardiovaskular pada Pasien Hipertensi dengan Risiko Tinggi: Laporan Berbasis Bukti

Alvin Nursalim,* Yoga Yuniadi#

Latar belakang: Pasien hipertensi dengan risiko tinggi mempunyai kemungkinan lebih tinggi mendapatkan komplikasi kardiovaskular. Pada pasien hipertensi risiko tinggi lebih baik digunakan obat hipertensi yang memiliki efek kardioprotektif. Angiotensin II Receptor Blocker (ARB) adalah salah satu golongan obat hipertensi yang memiliki efek kardioprotektif pada pasien hipertensi tanpa risiko. Apakah ARB juga efektif untuk menurunkan kejadian kardiovaskular baru atau perburukan kardiovaskular pada populasi hipertensi dengan risiko tinggi?

Tujuan: Mengetahui efektivitas ARB dibandingkan non-ARB dalam menurunkan komplikasi kardiovaskular pada pasien hipertensi dengan risiko tinggi.

Metode: Pencarian terstruktur dilakukan menggunakan Pubmed dan Cochrane. Setelah dilakukan penapisan judul dan abstrak, Tiga studi ditemukan yang kemudian digunakan penulis. Studi ini ditelaah dengan menggunakan kriteria yang mencakup *validity, importance, dan applicability* untuk menentukan derajat kegunaan studi.

Hasil: Berdasarkan studi oleh Sawada et al, valsartan memiliki efek penurunan tekanan darah sebanding dengan pengobatan non-ARB. Komplikasi kardiovaskular lebih rendah pada kelompok valsartan dibandingkan kelompok non-ARB (relative risk: 0.54, 95% confidence interval 0.4-0.7, $p < 0.001$). Valsartan menurunkan kejadian angina pectoris dibandingkan grup non-ARB (relative risk : 0.52, 95% Confidence Interval 0.31–0.86, $P = 0.01058$). Cohn JN et al menyimpulkan tidak ada perbedaan signifikan dalam pencegahan mortalitas dan parameter akhir lainnya. Namun penggunaan candesartan secara signifikan mengurangi kejadian diabetes.

Kesimpulan: Valsartan, dibandingkan pengobatan non-ARB, menurunkan kejadian kardiovaskular pada pasien hipertensi dengan risiko tinggi.

(J Kardiol Indones. 2012;33:91-8)

Kata kunci: Angiotensin II receptor blocker, valsartan, hipertensi risiko tinggi, komplikasi kardiovaskular

Alamat Korespondensi:

Dr. dr. Yoga Yuniadi, SpJP. Departemen Kardiologi dan Kedokteran Vaskular FKUI, dan Pusat Jantung Nasional Harapan Kita, Jln. S Parman Kav 87 Slipi Jakarta 11420. E-mail: yogayun@yahoo.com

Clinical scenario

A 58 years old man with dyslipidemia and type 2 diabetes mellitus diagnosed one year ago, came to the outpatient clinic for a regular check up in a private hospital. His physical examination was within

normal limit except for his high blood pressure (150/90). He was taking an ACE-inhibitors drug for his hypertension, but he complained of coughing not long after taking the drug. Due to the high incidence of cardiovascular complication among diabetic hypertensive patients, the doctor was thinking of changing his hypertensive drugs regimen to maximize the cardioprotective effect of the drug. The doctor wonders whether the administration of ARB, instead of non-ARBs, is more effective in preventing new onset or worsening of cardiac complication in high risk hypertensive patients.

Introduction

It is well established that hypertension is a cardiovascular disease risk factor. Hypertension also increases the risk of atherosclerosis progression.¹ Currently, the main cardiovascular disorder responsible for the rise in mortality has changed. As the substitute of rheumatic heart disease, atherosclerotic is now the leading cause for cardiovascular mortality. It is often assumed that atherosclerosis is a disease of high-income, industrialized countries. However, 80% of cardiovascular mortality occur in low-to-middle income countries.² Therefore, Indonesia as a developing country pose a high risk of having a high cardiovascular mortality.

High risk hypertensive patients will have an accelerated rate of atherosclerotic progression. In this report we try to determine the cardioprotective property of ARBs in high risk hypertensive patients (defined by patients with type 2 diabetes mellitus, lipid metabolism abnormality, obesity, and heart failure). Dyslipidemia is a strong risk predictor for coronary artery disease (CAD)³. Diabetic patients have twice to four times the risk of developing CAD as compared to those who do not have diabetes. Concomitant hypertension

triples the already high risk of CAD, doubles total mortality and stroke risk.⁴

Angiotensin II Receptor Blockers (ARB) have protective clinical effects similar to those of Angiotensin Converting Enzyme Inhibitors (ACE-Is) but better tolerated due to less side effect.⁵ Many studies have investigated the comparison of ARB and non-ARBs for stroke prevention, cardiac protection and renal protection.^{6,7} However, the efficacy of ARB as compared to non-ARBs to prevent new onset or worsening of cardiac complication on a more specific population, such as high risk patients need to be further elaborated.

The wide arrays of antihypertensive drugs available and the risk factors differences of each hypertensive patients made it tricky to choose the best anti-hypertensive regimen to reduce cardiac complication for these patients. This report is made to assist physician and investigate the efficacy of ARB compared to non-ARBs medication to prevent new onset or worsening of cardiac complication in high risk hypertensive patients.

Clinical question

Is ARB more effective compared to non-ARBs in preventing cardiac event in high risk hypertensive patients?

Methods

Search Strategy

The search was conducted on PubMed,[®] and Cochrane,[®] on February 8th 2011, using the keywords “angiotensin receptor blocker”, “hypertension” and

Table 1. Search strategy used in PubMed and Google (Conducted on November 11th 2011)

Database	Search terms	Results
Pubmed (8 th February 2011)	ARB[All Fields] AND (“cardiovascular system”[MeSH Terms] OR (“cardiovascular”[All Fields] AND “system”[All Fields]) OR “cardiovascular system”[All Fields] OR “cardiovascular”[All Fields]) AND complication[All Fields] AND (“epidemiology”[Subheading] OR “epidemiology”[All Fields] OR “morbidity”[All Fields] OR “morbidity”[MeSH Terms])	7
Cochrane (8 th February 2011)	“Hypertension” and “ARB” and “cardiovascular complication”	26

“cardiovascular complication” along with its synonyms and related terms (Table 1). Search strategy, results, the inclusion and exclusion criteria are shown in a flowchart (Figure 1).

Selection

The first selection was based on title and abstract using inclusion criteria and exclusion




Figure 1. Flow Chart of Search Strategy

criteria. After selection three full-text articles were available.

Critical Appraisal

After the selection, critical appraisal was done using several aspects based on Center of Evidence-based Medicine, University of Oxford for therapy study (Table 2).

preventing morbidity and mortality in high risk hypertensive patients. The study was a multicentre, prospective, randomized, open-label, blinded-endpoint trial. A total of 3.031 high risk hypertensive patients were randomized to either valsartan add-on or non-ARB treatment (ACE-Inhibitor was excluded). Patients with one or more of these risk factors is included: type 2 diabetes mellitus, current smoking, lipid metabolism abnormality, obesity (defined as BMI >25 kg/m²), and left ventricular hypertrophy defined

Table 2. Critical Appraisal of thesefull articles based on criterias by Centre of Evidence Medicine University of Oxford¹¹

Articles	Validity							Relevance			Result	Levels of evidence *
	Study design	Number of patients	Randomization	Similarity treatment and control	Blinding	Comparable treatment	Intention to treat	Domain	Determinant	Measurement of outcome		
Sawada et al ⁸	+	3,301	+	+	+	+	+	+	+	+	A	1B
Kasanuki et al ⁹	+	2049	+	+	+	+	?	+	+	+	B	1B
Cohn J N et al ¹⁰	+	5010	+	+	+	+	?	-	+	+	C	1B

Legend: + stated clearly in the article
 - not being done
 ? not stated clearly

*Levels of evidence based on The Oxford Centre of Evidence-based Medicine
 Domain: High risk hypertensive patients
 Determinant: Valsartan add-on therapy, Comparison: non-ARB therapy
 Outcome: New onset and/or worsening of cardiovascular event.
 A: The cardiovascular events in valsartan group was lower compared to non-ARB groups (relative risk: 0.54, 95% confidence interval 0.4-0.7, p< 0.001).
 B: There was no significant differences in the number of total death and primary endpoints. The only significant finding in this study is the occurrence of diabetes mellitus (Hazard ratio: 0.37; 95% CI, 0.16–0.89; P= 0.03).
 C: The rate of mortality was similar in the two treatment groups. The combined endpoint of mortality and morbidity was significantly reduced among patients in the valsartan group as compared to those in the placebo group (relative risk: 0.87, 95% confidence interval 0.77-0.97, p= 0.009).

Result

Sawada T et al⁸ performed a study to determined the efficacy of valsartan as compared to non ARBin

by electrocardiogram. The endpoints to be analyzed include new onset or worsening cardiovascular events, cerebrovascular events, mortality, worsening of cardiac function, new occurrence or exacerbation

of arrhythmias and new occurrence or exacerbation of diabetes mellitus.

The blood pressure lowering effect was similar between valsartan and non-ARB groups. The cardiovascular events was 83 out of 1517 and 155 out of 1514 in valsartan and non-ARB groups respectively (Hazard ratio: 0.55, 95% confidence interval 0.4-0.7, $P=0.00001$). The Control Event Rate (CER) was 10.2%, Experimental Event Rate (EER) was 5.5%, Relative Risk Reduction (RRR) was 46%, Absolute Risk Reduction (ARR) was 4.7% and Number Needed to Treat (NNT) was 21.

Another significant outcome in this study was the occurrence of angina pectoris. The administration of valsartan was superior to non-ARB in reducing angina pectoris. 22 patients given valsartan had angina pectoris compared with 44 in non-ARB group. The CER was 2.9%, EER was 1.5%, RRR was 48%, ARR was 1.4% and NNT was 71 (Hazard ratio: 0.51, 95% Confidence Interval 0.3-0.9, $P=0.01058$).

Cerebrovascular event represented by the occurrence of stroke also showed a significant reduction in the valsartan group compared to non-ARB group (Hazard ratio: 0.55, 95% Confidence Interval 0.3-0.9, $P=0.01488$).

Kasanuki H et al⁹ conducted a study to evaluate the efficacy of candesartan administration in reducing the incidence of cardiovascular events compared with non-ARB therapy in coronary artery disease patients with hypertension. A total of 2,049 patients were randomized to either candesartan therapy or non-ARB therapy (ACE-inhibitor was included).

The result of this study showed that there was no significant differences in the number of total death and primary endpoints, which include: cardiovascular deaths, non fatal myocardial infarction and heart failure. The only significant finding in this study is the occurrence of diabetes mellitus (hazard ratio: 0.37; 95% CI, 0.16-0.89; $P=0.03$).

Cohn J L et al¹⁰ evaluated the long term effects of the addition of valsartan to standard therapy for heart failure. This study was a randomized, placebo-controlled, double blind clinical trial. This is a large study involving 5,010 patients from 302 centers in 16 countries. Patients with heart failure were randomly assigned to receive 160 mg of valsartan or placebo twice daily. The primary endpoint was mortality and morbidity.

The result of this study showed that the rate of mortality was similar in the two treatment groups.

The combined endpoint of mortality and morbidity (hospitalization, cardiac arrest with resuscitation and intravenous therapy) was significantly reduced among patients in the valsartan group as compared to those in the placebo group. There were 723 patients of cases in the valsartan group and 801 cases in the placebo group (relative risk: 0.87, 95% confidence interval 0.77-0.97, $p=0.009$). The value of CER was 32%, EER was 29%, RRR was 9.4%, ARR was 3% and NNT was 33.

Discussion

Sawada et al⁸ studied the effect of ARB add on therapy on mortality and morbidity as compared to non-ARB treatment. Besides cardiovascular complication, the endpoint also includes cerebrovascular event, mortality and new onset diabetes. This study showed that the administration of valsartan therapy reduces the occurrence of cardiovascular event as compared to non-ARB therapy. The incidence of cardiovascular events was 5.5% and 10.2% in valsartan add-on and non-ARB groups, respectively. The relative risk reduction was 46% in the valsartan group as compared to the non-ARB group. There was also a lower rate of angina pectoris in the valsartan group as compared to non-ARB group (RRR=48%). The administration of valsartan was considered safe due to minimal adverse effects found throughout the study. The NNT for cardiovascular prevention was 21, making valsartan a reasonable drug to be applied widely.

The study by Kasanuki H showed different results.⁹ According to this study there was no significant difference in cardiovascular events between the candesartan group and placebo group. This conflicting result might result from some limitations of the study which include the high usage (71%) of ACE-inhibitors in the non-ARB therapy group and the low dose of candesartan used in the ARB group. However, the occurrence of diabetes was lower in the candesartan group. This protective effect of candesartan on diabetes supports the result of the VALUE study¹². According to this study ARBs were more effective in preventing diabetes compared to CCB.

The cardioprotective effect of valsartan is further ascertained in another study on valsartan performed by Cohn JN¹⁰. The third article by Cohn JN et al might have different patient characteristics as compared to our clinical question and the first two studies. This article studied the efficacy of valsartan as compared

to placebo for mortality and morbidity prevention among heart failure patients. Nevertheless, this study provide important information as addition to the first two studies, so we decided to include this study to the review. The result of this large scale study showed that valsartan was superior in reducing combined endpoint of mortality and morbidity (hospitalization, cardiac arrest with resuscitation and intravenous therapy) in patients with heart failure. Despite the high dose of valsartan (160 mg) used in this study, the drug was well tolerated and side effects were only slightly more prevalent in valsartan group. We do think the efficacy of valsartan still can be applied to hypertensive patients with heart failure. That is, under the assumption that the beneficial effects of valsartan among heart failure patients also apply to hypertensive patients with heart failure.

ARBs are well tolerated due to minimal side effects but the next intriguing question would be whether ARBs are as effective as ACE-Inhibitor in preventing cardiac events among high risk individual. According to two large scale studies (VALIANT study and ONTARGET study), ARBs are as effective as ACE-inhibitors for cardiovascular endpoints among high risk patients. This primary endpoint include cardiac related mortality, myocard infarct and stroke.^{13,14}

The administration of ARBs, in this case valsartan and candesartan, is considered safe due to the minimal side effect occurred during the study. Beside efficacy and safety, one aspect to consider when prescribing drug in Indonesia is the price of the drug. Currently, with "out of pocket" payment system (instead of health care insurance) and the low economic background of some patients, made it prudent for clinician to always consider drug cost in every clinical decision. ARBs are usually more expensive than other hypertensive drugs, fortunately some of these drugs are now available as generic drugs. It would be wise for clinician to individually tailored every drug prescription according to patients clinical condition and socio-economic background.

One particular limitation of this report might related to the search strategy. In terms of the search strategy, we admit that we might have missed relevant articles due to the exclusion of all articles that were not available with full text on the internet. We only found two articles which perfectly related to our clinical question. Nonetheless both articles are large scale studies with adequate follow-up period and they reported similar benefit of ARBs (valsartan and

candesartan). Clinical experience with these ARBs are quite extensive, for this reason ARBs can be considered as a cardioprotective hypertensive agent for high risk hypertensive patients.

Conclusion and recommendation

ARB therapy, represented by valsartan and candesartan in these studies, is effective in reducing cardiovascular event in high risk hypertensive patients (defined by patients with type 2 diabetes mellitus, lipid metabolism abnormality, obesity, and heart failure). Valsartan therapy is well tolerated, as shown by the minimum adverse effects found throughout these studies. ARBs tend to be more expensive, therefore, the best implementation of this evidence requires risk factors and socio-economic assessment of each patient.

This new insight can be translated into clinical practice. Based on this evidence, we recommend the administration of valsartan, an ARB drug, for high risk hypertensive patient in the clinical scenario mentioned above.

References

1. Alexander RW. Hypertension and the Pathogenesis of Atherosclerosis Oxidative Stress and the Mediation of Arterial Inflammatory Response: A New Perspective. *Hypertension*. 1995;25:155-61
2. Bonow RO, Smaha LA, Smith SC, Mensah GA, Lefant C. World Heart Day 2002: The International Burden of Cardiovascular Disease: Responding to the Emerging Global Epidemic. *Circulation* 2002, 106:1602-05.
3. Kavey REW, Daniels SR, Lauer RM, Atkins DL, Hayman LL and Taubert K. American heart Association Guidelines for Primary Prevention of Atherosclerotic Cardiovascular Disease Beginning in Childhood. *Circulation* 2003, 107:1562-66.
4. Stults B and Jones RE. Management of Hypertension in Diabetes. *Diabetes Spectrum*, 2006, 19(1):25-31.
5. Birkenhäger WH, de Leeuw PW. Non-peptide Angiotensin type 1 Receptor Antagonists in The Treatment of Hypertension. *J Hypertens* 1999;17:873-81.
6. Wang JG, Li Y, Franklin SS, Safar M. Prevention of Stroke and Myocardial Infarction by Amlodipine and Angiotensin Receptor Blockers: A Quantitative Overview. *Hypertension* 2007;50:181-8.
7. Pfeffer MA, McMurray JJV, Velasquez EJ, Rouleau JL, Kober

- L, Maggioni AP et al. Valsartan, Captopril, or Both in Myocardial Infarction Complicated by Heart Failure, Left Ventricular Dysfunction, or Both. *N Eng J Med* 2003;349:1893-1906.
8. Sawada T, Yamada H, Dahlof B, Matsubara H. Effects of Valsartan on Morbidity and Mortality in Uncontrolled Hypertensive Patients with High Cardiovascular Risk: Kyoto Heart Study. *Eur Heart Journal* 2009; 30: 2461-69.
 9. Kasanuki H, Hagiwara N, Hosada S, Sumiyoshi T, Honda T, Haze K et al. Angiotensin II receptor blocker-based vs non-angiotensin II receptor blocker-based therapy in patients with angiographically documented coronary artery disease and hypertension: the Heart Institute of Japan Candesartan Randomized Trial for Evaluation in Coronary Artery Disease (HIJ-CREATE). *European Heart Journal* 2009; 30: 1203-12.
 10. Cohn JN, Tognoni G. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med* 2001; 345:1667-75.
 11. Centre for Evidence Based Medicine. Oxford Centre for Evidence-based Medicine- Level of Evidence. CEBM. [cited 2012 February 5]. Available from: URL:HYPERLYNKhttp://www.cebm.net/index.aspx?o=5513
 12. Julius S, Weber MA, Kjeldsen SE, McInnes GT, Zanchetti A, Brunner HR et al. The valsartan antihypertensive long-term use evaluation (VALUE) trial: outcomes in patients receiving monotherapy. *Hypertension* 2006; 48:385-91)
 13. The ONTARGET Investigators. Telmisartan, Ramipril, or Both in Patients at High Risk for Vascular Events. *N Eng J Med* 2008; 358:1547-59.
 14. Pfeffer MA, McMurray JJV, Velazquez EJ, Rouleau JL, Kober L, Maggioni AP et al. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Eng J med* 2003; 349:1893-906.